



LIVISTO

GANAMINOVIT

READY TO PLAY

Vitamin and aminoacid oral powder



Along with you

GANAMINOVIT

Vitamin and aminoacid oral powder



READY TO PLAY

It is well known that a deficiency in vitamins and amino acids in the diet may bring about a lot of problems in the farm. In the chickens, it may lead to:

- worse carcass quality
- muscular dystrophy
- neurologic lesions
- drop in egg production
- higher embryonic mortality
- death of the animal



The diet fed to the chicks in their first week of life is crucial to determine the final weight of the chicken.

GANAMINOVIT contains all the liposoluble vitamins (A, D, E, and K₃) and the most important water soluble vitamins (B₁, B₂, B₃, B₅, B₆, biotin, folic acid, B₁₂ and C). Besides, it has all the essential amino acids and a great variety of non-essential amino acids.

Provide all necessary nutrients in the farm to ensure the health and the optimal performance of the animals with GANAMINOVIT.

HOW TO ADMINISTER GANAMINOVIT?

Mix 1 kg of GANAMINOVIT per 1000 L of water, for 4-8 days.
*Medicated water should be used within 24h.

Composition 1 g contains: Vitamin A, 10,000 IU; Vitamin D₃, 1,000 IU; Vitamin E, 10 mg; Vitamin B₁ hydrochloride, 2 mg; Vitamin B₂, 4 mg; Calcium pantothenate, 10 mg; Vitamin B₆ hydrochloride, 1.5 mg; Vitamin C, 25 mg; Vitamin K₃, 1.5 mg; Folic acid, 500 µg; Nicotinamide, 20 mg; Biotin, 15 µg; DL-Methionine, 50 mg; L-Lysine, 50 mg; Hydrolysed vegetable proteins (which provides: Glutamic acid, Proline, Leucine, Aspartic acid, Valine, Serine, Isoleucine, Phenylalanine, Threonine, Arginine, Alanine, Histidine, Glycine, Tryptophan and Cysteine), 250 mg. **Indications and target species** Poultry, cattle, sheep, goats and horses: Stress conditions, improper feed, poor condition. **Contraindications, adverse reactions and interactions** Not described. **Dosage for each species, route and method of administration** Oral use in drinking water. Poultry: 1-2 g/2 L of drinking water for 4-8 consecutive days. Cattle, sheep, goats and horses: 1 g/L of drinking water for 3-5 consecutive days. It is recommended to make a presolution in about 10 L of water. Medicated drinking water should be used within 24 hours. **Withdrawal period** Not necessary. **Storage precautions** Keep out of the sight and reach of children. Protect from moisture, light and heat. **Other information** Containers of 100 g, 1 kg and 25 kg. Not all pack sizes may be marketed. **For animal use.**