



ANIMELOXAN

20 mg/ml

DON'T LET THE FLAME BURN

Meloxicam solution for injection for cattle, pigs and horses

LESS GASTROINTESTINAL EFFECTS. SINGLE DOSE

ANIMELOXAN 20 mg/ml solution for injection for cattle, pigs and horses. Composition: One ml contains: Meloxicam 20 mg. **Target species:** Cattle, pigs and horses. **Contraindications:** Do not use in horses less than 6 weeks of age. Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions. Do not use in case of hypersensitivity to the active substance or to any of the excipients. For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age. **Special precautions for use:** **Special precautions for use in animals:** If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought. Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity. In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention. **Special precautions to be taken by the person administering the veterinary medicinal product to animals:** Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician. If accidental skin contact occurs, wash the affected area thoroughly. Wash hands after use. In view of the risk of accidental self-injection and the known adverse class-effects of NSAIDs and other prostaglandin inhibitors on pregnancy and/or embryofetal development, the veterinary medicinal product should not be administered by pregnant women or women attempting to conceive. **Withdrawal periods:** Cattle: Meat and offal: 15 days; Milk: 5 days. Pigs: Meat and offal: 8 days. Horses: Meat and offal: 5 days. Do not use in horses producing milk for human consumption. **Special precautions for storage:** This veterinary medicinal product does not require any special temperature storage conditions. **Packaging:** Glass vials of 50 ml and 100 ml. **Marketing Authorisation Holder:** aniMedica GmbH, Im Südfeld 9, 48308 Senden-Bösensell, Germany.



Along with you

livisto.com