



SURRICOXX[®] 400 mg/mL

Solution for use in drinking water

Contains per mL: Amprolium 400.0 mg (equivalent to 452.4 mg of amprolium hydrochloride)



of Amprolium



- ✓ For chickens, turkeys, ducks and guinea fowls
- ✓ Highly concentrated
- ✓ Withdrawal time: 0 days for eggs, meat and offalls

SURRICOXX[®] 400 mg/mL

Solution for use in drinking water

Contains per mL: Amprolium 400.0 mg (equivalent to 452.4 mg of amprolium hydrochloride)




Indications

Treatment of intestinal coccidiosis caused by *Eimeria* spp. susceptible to amprolium.



Contraindications

Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.



Adverse reactions

None



Withdrawal period

- **Meat and offal:** zero days
- **Eggs:** zero days

Storage

Do not store above 30 °C.

Shelf life

- Shelf life of the veterinary medicinal product as packaged for sale: 21 months.
- Shelf life after first opening the immediate packaging: 3 months.
- Shelf life after dilution according to directions: 24 hours.

Presentation

100 mL, 1 L, 5 L

The impact on poultry production

- ✓ Worldwide, coccidiosis is one of the main diseases in the poultry sector*
- 6 to 10 % of mortality worldwide
- 5 % of broiler flocks are suffering from acute outbreaks
- 20 to 30 % of farms have subclinical infectious pressure

6 means of control

- ✓ Hygiene
- ✓ Zootechny
- ✓ Prevention with coccidiostat (Ionophore/chemical)
- ✓ Prevention with plant extracts
- ✓ Vaccination
- ✓ Treatment

Sources

- * The value of anticoccidials for sustainable global poultry production (Review); Stefanie Kadykalo, Tara Roberts, Michelle Thompson, Jeff Wilson, Marcelo Lang, Olivier Espeisse, International Journal of Antimicrobial Agents; Volume 51, Issue 3, March 2018, Pages 304-310
- * Anticoccidial drugs of the livestock industry; Sandra Noack, H. David Chapman, and Paul M. Selzer, Parasitol Res. 2019; 118(7): 2009–2026.

Dosage and administration

In drinking water.

for each target species	
20 mg of amprolium / kg body weight / day	
= 0.5 mL Surricoxx [®] / 10 kg body weight / day or 50 mL / T body weight / day	
for 5 to 7 consecutive days	

for the treatment of a flock					
0.05 mL of Surricoxx [®] / kg body weight	x	average body weight (kg) of the animals	x	number of animals	= mL of oral solution / L of drinking water, to be prepared each day
0.05 mL	x	... kg	xmL of Surricoxx [®]
Example: 10 000 chickens, each weighing 750 g, will receive 375 mL of Surricoxx [®] each day, for 5 to 7 days					

Solubility in drinking water of up to 100 mL product per litre can be used when preparing stock solutions for use in a water proportioner system which subsequently dilutes the product to its correct final concentration.